

BEYOND POPULAR SCIENCE



DAVID H. SILVER



BEYOND POPULAR SCIENCE

David H. Silver

<https://www.openbookpublishers.com>

© 2026 David H. Silver



This work is licensed under the Creative Commons Attribution-NonCommercial 4.0 International (CC BY-NC 4.0). This license allows you to share, copy, distribute and transmit the text; to adapt the text for non-commercial purposes of the text providing attribution is made to the authors (but not in any way that suggests that they endorse you or your use of the work). Attribution should include the following information:

David H. Silver, *Beyond Popular Science*. Cambridge, UK: Open Book Publishers, 2026,
<https://doi.org/10.11647/OBP.0526>

Further details about CC BY-NC licenses are available at
<https://creativecommons.org/licenses/by-nc/4.0/>

Copyright and permissions for the reuse of many of the images included in this publication differ from the above. This information is provided in the captions and in the list of illustrations. Unless otherwise stated, figures are reproduced under the fair dealing principle. Every effort has been made to identify and contact copyright holders and any omission or error will be corrected if notification is made to the publisher.

All external links were active at the time of publication unless otherwise stated and have been archived via the Internet Archive Wayback Machine at
<https://archive.org/web>

Digital material and resources associated with this volume are available at
<https://doi.org/10.11647/OBP.0526#resources>

ISBN Paperback:	978-1-80511-877-0
ISBN Hardback:	978-1-80511-878-7
ISBN Digital (PDF):	978-1-80511-879-4
ISBN HTML:	978-1-80511-881-7
ISBN Digital ebook (epub):	978-1-80511-880-0
DOI:	10.11647/OBP.0526




Cover image by Enny Silver and David H. Silver
Cover design by Jeevanjot Kaur Nagpal

A Place at the End of Time

Top (Spacetime Geometry): The geometry near a black hole distorts spacetime so severely that, from a coordinate perspective, the black hole itself lies in the future light cone of any nearby event. For an infalling observer, continuing forward in time inevitably means moving closer to the singularity—tomorrow *is* the black hole in spacetime terms.

Middle (Observer Perspectives): Two perspectives on crossing the horizon. From the distant observer's viewpoint (left), the astronaut appears to freeze at the horizon, increasingly redshifted and dimmed, never quite crossing. From the astronaut's own frame (right), there's no discontinuity—no freezing, no slowdown—only smooth, uninterrupted free fall through the horizon in finite proper time.

Bottom (Penrose Diagram): The full Penrose diagram for the maximally extended Schwarzschild solution, showing all causal regions: the external universe, black hole interior, white hole region, and a second asymptotically flat universe. The diagram captures the global causal structure: horizons, infinities and singularities.



A Place at the End of Time

Black holes create an observational paradox: external observers see infalling objects freeze at the event horizon with infinite redshift, while the falling objects cross in finite proper time experiencing nothing unusual. This contradiction arises from extreme spacetime curvature near the horizon ($r = 2GM/c^2$), where gravitational time dilation becomes unbounded. Inside the horizon, causality inverts—the radial coordinate becomes timelike, making the singularity not a place but a future moment that all trajectories must reach.



BLACK HOLE EVENT HORIZON ◦ SCHWARZSCHILD
RADIUS ◦ TIME BECOMES SPACELIKE ◦ SINGULARITY AS
FUTURE ◦ LIGO GRAVITATIONAL WAVES ◦ EHT M87
IMAGE ◦ COORDINATE DEPENDENCIES ◦ PENROSE-HAWKING
THEOREMS ◦ KERR'S QUORA OBJECTIONS ◦ WHITE HOLES &
WORMHOLES ◦ r BECOMES TIMELIKE

“So much universe, and so little time.”

— Cohen the Barbarian, AM 1980s

A Place at the End of Time

Karl Schwarzschild derived the first exact solution to Einstein's field equations in 1916, giving rise to what is now known as the Schwarzschild metric. This solution described the spacetime geometry outside a static, spherically symmetric mass, and revealed an intriguing radius where the metric becomes singular—a mathematical curiosity at the time.

In 1939, Robert Oppenheimer and Hartland Snyder explored the gravitational collapse of massive stars showing that under general relativity, such collapse could lead to the formation of a region from which no signals escape: the conceptual precursor to what we now call a black hole.

The term 'black hole' was popularised by John Wheeler in 1967, highlighting that these regions are not conventional objects, but causal domains shaped by the warping of spacetime. In 1963, Roy Kerr discovered an exact solution for rotating black holes. The Kerr metric demonstrated that black holes can possess angular momentum, vastly enriching the theory's physical relevance. Unlike Schwarzschild's static solution, the Kerr geometry features an ergosphere, frame dragging, and a more rich causal structure—including an inner and outer horizon.

Throughout the twentieth century, indirect astrophysical evidence for black holes mounted, from X-ray binaries to quasars and galactic nuclei. By the 2010s, observations of gravitational waves and the first black hole shadow image (captured by the Event Horizon Telescope in 2019) solidified their status as real astrophysical objects.

Despite this progress, black holes continue to raise deep questions. At their core lies the unresolved issue of singularities—regions where classical spacetime is undefined—and the challenge of unifying general relativity with quantum theory remains a central frontier in physics.

General relativity describes gravity as spacetime curvature. Massive bodies distort the geometry in which other bodies move, and free-fall corresponds to inertial motion along geodesics—paths of extremal proper time. This reconceptualization allows for solutions to Einstein's equations that have no Newtonian counterpart. When matter collapses to a sufficiently small region, the curvature becomes extreme enough that no causal signal can propagate outward beyond a critical boundary.

A black hole is defined by geometry. A region of spacetime in which spatial and temporal concepts blend. The defining feature is the event horizon: a null surface that separates regions of spacetime into two domains: those from which future-directed paths can reach infinity, and those from which all such paths terminate inward. The horizon has no surface tension or material properties. Its existence follows purely from the metric. In the Schwarzschild solution, the horizon forms (Schwarzschild, 1916) at radius $r = 2GM/c^2$, where the g_{00} component vanishes and light cones tip inward. Any trajectory, regardless of force or energy, once inside this radius, proceeds inevitably toward smaller r .

Such configurations are predicted results of stellar evolution. When a sufficiently massive star exhausts its nuclear fuel, no internal pressure, thermal, degeneracy, or radiation can

oppose further collapse. Neutron stars represent the final stable configuration for masses up to a few solar masses. Beyond that, collapse continues past any known state of matter. General relativity predicts that the outer region smooths into a vacuum solution matching Schwarzschild or Kerr metrics, while the interior (Kerr, 1963) forms a trapped surface with inward-pointing causal futures. The event horizon forms before any singularity becomes visible, preventing external observers from accessing information about the final collapse state.

This scenario was further confirmed in 2015—when LIGO detected gravitational waves from a binary black hole merger. The distortion in spacetime, measured to better than one part in 10^{21} , was generated by two orbiting black holes coalescing into one. The signal matched numerical relativity simulations, confirming the waveform, mass loss, and final ringdown predicted by general relativity. LIGO thus became one of the most sensitive measurement devices ever built, detecting strains comparable to changes smaller than a proton over kilometre-scale arms. The black holes radiated energy equivalent to several solar masses through measurable curvature oscillations.

Other confirmations have followed. The Event Horizon Telescope array imaged the shadow of the supermassive black hole in M87, producing a crescent-shaped brightness profile consistent with light bending and lensing near the photon sphere. Stellar orbit measurements around Sagittarius A* in the centre of the Milky Way reveal elliptical motions governed by a central mass of approximately four million solar masses in a region smaller than the orbits themselves. Accretion disc X-ray emissions, variability timing, and iron line broadening all support the interpretation of compact objects with deep gravitational wells: exhibiting effects that match the metrics of rotating (Kerr) black holes with no observable surface.

As one approaches a black hole, time ceases to behave as expected. The component g_{00} of the spacetime metric determines how proper time accumulates for a stationary observer. In Schwarzschild geometry, $g_{00} = 1 - 2GM/rc^2$ decreases with decreasing radius. A clock closer to the event horizon ticks more slowly relative to one farther away. The gravitational redshift of light signals this disparity—photons emitted near the horizon lose energy as their wavelengths stretch. At the horizon, the redshift becomes unbounded. Signals emitted at or within the horizon do not reach distant observers; emissions from just outside arrive with arbitrarily large delay and redshift.

An object falling into the black hole measures finite proper time to cross the event horizon; locally, nothing singular occurs there (neglecting tidal forces). This dual description, freezing from the outside, flowing from the inside, follows from the coordinate-dependence of simultaneity in general relativity. Infalling observers describe the event horizon as a regular null surface. The difference lies in the slicing of spacetime used to define simultaneity. Proper time and coordinate time diverge in meaning as curvature intensifies.

Inside a black hole (in Schwarzschild coordinates), the radial coordinate behaves time-like—decreasing radius corresponds to forward progression in time—while the temporal coordinate behaves spacelike. In horizon-regular coordinates, this role-swap is recognised as a coordinate effect; causality still directs all future paths toward smaller r .

Outside the horizon, the singularity occupies the spatial point $r = 0$. Inside, it changes from a place to a moment. The question is not ‘where is the singularity?’ but ‘when will I reach it?’ The answer: finite proper time ahead, as inevitable as tomorrow. Light cones inside the horizon all tilt toward smaller r , making motion toward the singularity as compulsory as motion into the future. Remaining at fixed radius would require stopping time.

The Penrose–Hawking theorems show that under reasonable energy (Penrose, 1965) and global conditions, spacetimes containing trapped surfaces are geodesically incomplete: certain timelike or null geodesics cannot be extended to arbitrary values of their affine parameter. This geodesic incompleteness—what is meant by a ‘singularity’ in this context—lies in the future of every worldline that crosses the event horizon in the idealised solutions. The theorems do not by themselves guarantee curvature divergence everywhere; rather, they establish the existence of incomplete causal paths. Thus, you cannot point to the singularity; it is a when, not a where.

The field equations of general relativity are time-symmetric. If the Schwarzschild solution describes an object into which signals can enter but never leave, then its time-reversed counterpart also exists. This reversed solution is called a white hole: a region of spacetime from which causal trajectories can emerge, but into which nothing can be sent. Unlike black holes, white holes cannot be formed dynamically under known physical processes. They appear in maximal analytic extensions (such as Kruskal spacetime) but lack known mechanisms for creation or stability.

Another extension is the wormhole: a spacetime manifold that connects two asymptotically flat regions through a throat. In its simplest form, the Einstein–Rosen bridge arises from a slicing (Einstein & Rosen, 1935) of the maximally extended Schwarzschild geometry. However, the bridge pinches off too rapidly to allow traversal. For a wormhole to be traversable, the geometry must remain open long enough for causal passage. This requires exotic matter: fields or fluids that violate the null energy condition, allowing repulsive gravitational effects. Such matter has not been observed. Moreover, semiclassical analyses suggest instabilities that would disrupt the throat, collapse the tunnel, or generate divergent backreaction.

Black holes, white holes, and wormholes demonstrate surprising configurations of spacetime. Near a black hole, coordinate roles switch, light cones tilt, and the metric enforces trajectories independent of any force. White holes, if they exist, have always existed. Wormholes in general relativity such as the Einstein–Rosen bridge are non-traversable; traversable wormholes would require exotic matter and remain hypothetical.

Kerr’s Quora Posts: “Stop Believing Everything You Read About Black Holes”

Sixty years after discovering the metric that bears his name, Roy Kerr has taken to Quora with the fury of a physicist whose life's work has been misinterpreted. His posts read like manifestos from an exile returning to reclaim his territory. ‘Stop believing everything you read about black holes,’ he declares, targeting not just popular misconceptions but the physics community.

Kerr's central accusation is that the Penrose singularity theorems prove nothing about physical singularities. What Penrose actually showed was that certain geodesics have finite affine length—they simply end. ‘Now, what if the central star is singular?’ Kerr asks pointedly. ‘Then one is assuming it is singular and there is nothing to prove.’ He claims circular reasoning: assume a singularity exists, then ‘prove’ singularities must exist.

His technical objection cuts directly to the heart of black hole physics. In the Kerr solution, geodesics starting outside can pass through a central neutron star and terminate on the inner horizon on the opposite side. These are Penrose's ‘mysterious light rays of finite affine length.’ They die not because they hit an infinitely curved singularity, but because they complete their journey through the black hole's interior. Geodesic incompleteness is a boundary condition rather than a catastrophe.

The medium amplifies the message. Quora allows Kerr to bypass peer review and speak directly: ‘The trouble with the Penrose paper is that it is a “do it yourself” paper where he states propositions without proving them. This is very typical in relativity [...] conjectures “rule the roost.”’ These are not the measured tones of academic discourse but the exasperated words of someone watching decades of what he considers misinterpretation compound.

Most provocatively, Kerr disputes the coordinate interpretation that underlies this entire chapter. Asked whether ‘time and space exchange roles at the event horizon,’ his response is unequivocal: ‘Of course this is not true.’ The coordinate swap reflects bad coordinate choices, not physics. ‘Time is a function defined on a physical manifold with the property that it increases along every world line,’ he explains, demolishing the temporal interpretation in a single stroke.

The technical details matter. In Kerr-Schild coordinates, which Kerr considers ‘good,’ the t -coordinate remains a proper time parameter along all worldlines, never becoming spacelike. The dramatic coordinate inversions described throughout this chapter— r becoming timelike, the singularity becoming temporal—are artefacts of choosing Schwarzschild coordinates, which produce a t -coordinate that ‘is not a differentiable function on the manifold.’ Use better coordinates, and the mystery vanishes.

Yet Kerr's alternative is equally radical. He describes ‘spin forces’ that become so intense near the event horizon that infalling objects are forced to rotate around the axis. At the inner horizon, centrifugal forces grow strong enough that objects can move outward again—no longer forced toward any central singularity.

The stakes are higher than academic priority. If Kerr is correct, then black holes are not the temporal futures described in this chapter but something else entirely: regions where extreme spin and gravity create exotic dynamics. His Quora posts are interesting not only for the physics or for the fact that he first solved the stable black hole equations, but also for the platform and direct style of his writing.

It's Ferret Time

A guy's car breaks down on a rural road. No cell phone reception. Fortunately, there's a farmhouse nearby, so he walks over to ask for help.

He rings the doorbell—no answer. But he hears rustling from around the side. He finds a farmer next to a pen with three ferrets.

'Excuse me—' the guy begins, but the farmer cuts in: *'Hold on, I have to feed the ferrets. I'll be with you when I'm done.'*

The guy watches as the farmer picks up one of the ferrets, carries it to an apple tree with a ladder, climbs up while holding it, lifts it to an apple—*chomp*—then climbs back down and returns it to the pen.

'Sorry, I just—' the guy tries again, but the farmer grabs the second ferret. *'Almost done.'*

Same thing: apple tree, ladder, apple, chomp, back down, pen.

Then the third ferret. The farmer lifts it gently, murmurs something to it, and makes the familiar journey. Step by step up the ladder, hoists the ferret to a fresh apple—another bite. He climbs down slowly, cradles it all the way back, nestles it in the pen like a newborn.

Finally, the farmer turns. *'Now—how can I help you?'*

The guy says, *'My car broke down and I need to call AAA. But first... wouldn't it be a lot faster to just pick the apples and toss them into the pen?'*

The farmer pauses, thinking. Then nods. *'Yeah... I suppose it would be faster that way.'*

He shrugs.

'But what's time to a ferret?'

Coordinate Reversal

Schwarzschild Metric and the Horizon

The Schwarzschild metric describes space-time outside a static mass M . In spherical coordinates (t, r, θ, ϕ) , it takes the form (with $r_s = 2GM/c^2$):

$$ds^2 = - \left(1 - \frac{r_s}{r}\right) c^2 dt^2 + \left(1 - \frac{r_s}{r}\right)^{-1} dr^2 + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2.$$

At $r = r_s$, $g_{tt} = 0$, $g_{rr} \rightarrow \infty$. This signals a breakdown in coordinates, not in geometry.

Time Dilation and Gravitational Redshift

A static observer at fixed radius $r > r_s$ experiences proper time:

$$d\tau = \sqrt{1 - \frac{r_s}{r}} dt.$$

As $r \rightarrow r_s$, $d\tau/dt \rightarrow 0$. Distant clocks appear to tick normally, but local clocks slow near the horizon.

Photons emitted at r_{em} and received at $r_{\text{obs}} \rightarrow \infty$ undergo redshift:

$$1 + z = \left(1 - \frac{r_s}{r_{\text{em}}}\right)^{-1/2}.$$

As $r_{\text{em}} \rightarrow r_s$, $z \rightarrow \infty$. Light from near the horizon becomes infinitely stretched and fades from view.

Finite Infall and Apparent Freezing

A freely falling observer released from rest at radius $r_0 > r_s$ reaches radius $r \leq r_0$ in proper time:

$$\tau(r; r_0) = \frac{2}{3} \frac{r_0^{3/2} - r^{3/2}}{\sqrt{2GM}}.$$

For any finite r_0 , the proper time to reach the horizon is finite. The infaller feels no discontinuity; the horizon is not a physical surface.

Even an infaller starting from rest at infinity crosses the horizon in finite proper time, and continues to $r = 0$ also in finite proper time.

However, to a distant observer, the infaller appears to freeze at $r = r_s$, due to the divergence of coordinate time: $t(r) \rightarrow \infty$ as $r \rightarrow r_s$.

Coordinate Inversion Below the Horizon

Inside the horizon ($r < r_s$), the signs of metric components reverse in Schwarzschild coordinates: $g_{tt} > 0$, $g_{rr} < 0$. In this coordinate choice r behaves timelike and t spacelike; in horizon-regular coordinates (e.g., Eddington–Finkelstein, Kruskal) this interpretation is seen as a coordinate effect while causal futures still point to decreasing r .

The physical implication is that movement in r becomes mandatory. All future-directed timelike paths lead to smaller r , ending at the singularity $r = 0$. The singularity is not ‘a place inside’ but a moment in the infaller’s proper future.

This inversion reflects the geometry. Any attempt to ‘hover’ or remain at constant r is no longer physically possible once inside the horizon.

Light Cones and Irreversibility

At the horizon, outgoing light rays remain on the surface: $\frac{dr}{dt} = 0$ for outgoing null rays at $r = r_s$.

Below the horizon, all light cones tip inward. Future lightlike and timelike paths are directed toward decreasing r . There is no direction within the cone that leads to increasing radius.

This defines the event horizon as a one-way temporal boundary: a surface from which causal influence cannot escape outward.

References:

- Misner, C. W., Thorne, K. S., Wheeler, J. A. (1973). *Gravitation*.
 Carroll, S. M. (2004). *Spacetime and Geometry*.
 Wald, R. M. (1984). *General Relativity*.

